

Hormonal and Reproductive Factors Associated with Low Back pain in Reproductive Female: A Hospital Based Study in Mathura

Aparajita Gulati, Nupur Ahuja*

Assistant Professor, (Obs and Gynae),
Krishna Mohan Medical College & Hospital,
Mathura

ABSTRACT

Background: Low back pain related disability and work absence accounts for high economical costs in modern society. A prevalence of 28-80% has been found with increase of prevalence with age and female preponderance.

Methods: The present study was conducted on 250 non pregnant women complaining of low back pain attending Department of Obstetrics and Gynaecology, during the period of one year. A brief questionnaire was used to screen, among the respondents, the occurrence of low back pain in the past year.

Results: Low back pain was associated with high BMI, high waist circumference, more number of children, irregular and prolonged duration of menstruation, young maternal age at first birth and with history of abortion.

Conclusions: Hormonal and reproductive factors are associated with low back pain.

Keywords: Hormonal factors, Reproductive factors, Reproductive female

Available Online: 30th June 2020

Received: 20.05.20

Accepted: 18.06.20

*Corresponding Author

Dr. Nupur Ahuja,
Assistant Professor, Obst. & Gynae,
Krishna Mohan Medical College & Hospital,
Pali Dungra, Sonkh Road, Mathura
Email: kmmch.nupur@gmail.com

Copyright: © the author(s). IABCR is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.




This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial

INTRODUCTION

Low back pain is defined as pain and discomfort, localized below the costal margin and above the inferior gluteal folds, with or without referred leg pain. Chronic low back pain is defined as low back pain persisting for at least 12 weeks, unless specified otherwise. A simplified and practical classification, which has gained international acceptance, is to divide low back pain into three categories-the so called "diagnostic triage"¹ : 1. Specific spinal pathology 2. Nerve root pain/radicular pain 3. Non-specific low back pain. Low back pain is common disorder, affecting around one-

third of UK adult population each year. About 20% of people of low back pain (that is 1 in 15 of population) will consult their GP about it. The presence of low back pain during pregnancy is widely reported.²⁻⁵ One systematic review identified 56 population prevalence studies of low back pain (Walker 2000). Thirty studies were of acceptable quality. Point prevalence of low back pain ranged from 12-33%. The two reviews on low back pain in school children and adolescents reported a prevalence approaching that

Access this article online	
Website: www.iabcr.org	Quick Response code 
DOI: 10.21276/iabcr.2020.6.2.16	

How to cite this article: Gulati A, Ahuja N. Hormonal and Reproductive Factors Associated with Low Back pain in Reproductive Female: A Hospital Based Study in Mathura. Int Arch BioMed Clin Res. 2020;6(2):OG4-OG6.

Source of Support: Nil, **Conflict of Interest:** None

reported for adults.^{6,7} Low back pain fluctuates over time with frequent recurrences and exacerbations.⁸ The first review reported that, after a first episode of low back pain, the proportion of patient who still experienced pain after 12 months was on average 62% (range 42-75%), the percentage who experienced relapses of pain was 60% (range 44-78%) and the percentage who had relapses of work absence was 33% (range 26-37%).⁹ Factors which are associated with low back pain are young age at menarche, irregular or prolonged menstruation, past pregnancy, young maternal age at first birth, and duration of oral contraceptive use, hysterectomy and use of estrogens during menopause.¹⁰ Other factors associated with low back pain are previous miscarriage, presence of endometriosis, clinically suspected pelvic inflammatory disease, caesarean section scar, pelvic adhesions, sexual abuse, anxiety, and depression. Back pain is a commonly described symptom of the premenstrual syndrome.¹¹ The aim and objectives of the study was to study the hormonal and reproductive factors associated with low back pain in reproductive female.

METHODS

The present study is a prospective study conducted on 500 subjects attending Department of Obstetrics and Gynaecology, over a period of 12 months. The subjects were randomly allocated and found 125 patients suffering from low back pain. Inclusion criterion 1. Women attending outpatient department of obstetrics and Gynecology in UISEMH. 2. All females of age 20-45 yrs of any religion and socioeconomic status. Exclusion criterion 1. Women who have attained physiological menopause. 2. Subjects who were not able to communicate because of dialect or hearing problems. Study protocol A brief questionnaire was used to screen, among the respondents, the occurrence of low back pain in the past year. The questions included occurrence of low back pain, demographic factors and reproductive health history. Low back pain was identified among subjects who have back pain lasting for more than a day in an area between the lower coastal margin and the gluteal folds with or without radiation into leg to below the knees during the past one year. The anthropometric measurements included measures of body height (cm) and weight (kg) using standard measurement equipment's. Waist and hip circumference (cm) were assessed using a measuring tape while the subject was standing. Other factors evaluated were occupation, education and smoking. Sociodemographic profile, menstrual and obstetric history were noted and also the use of oral contraceptives.

RESULTS

Out of 500 subjects, 125 were patients of low back pain. Majority of patients, 55.6% were in age range of 40-45 years and only 12.5% in 20-30 years of age group. Out of 125, 55.4% belongs to socioeconomic class 1 and 2. Only 36 patients had education of university and above. 24.8% (31/125) were Labourers and rest 75.2% had some other occupation. In our study 60% (75/125) patients had height below median (153 cm) whereas 39.6% (50/125) had their heights above median. Similarly, 55.2% (69/125) had weight below median (60 kg) and only 44.8% had weight above median.

Patients with BMI (>25kg/m²) were 55.6% and 44.6% had BMI below 25 kg/m². 55 out of 125 had waist circumference of <80 cm and 69 (59%) had more than 80 cm. Their waist-hip ratio is <0.8 in 65 patients (52.8%).

Majority of patients had more than two parity (64.8%). Patients with vaginal and caesarean delivery were 61 (48.8%) and 64 (51.2%) respectively. 72 patients out of 125 had irregular menstruation. Majority of them (64.8%) had duration of flow of more than 8 days. 82 patients had premenstrual syndrome. 69.2% patients had undergone sterilization and 41.2% (50/125) were using oral contraceptives. 80 patients had history of abortion and 38.8% patients had pelvic organ prolapse.

DISCUSSION

The study showed that hormonal and reproductive factors like irregular or prolonged menstrual cycle were associated with low back pain. Estrogen related factors like past pregnancy, young maternal age at first birth, oral contraceptive use was specifically associated with low back pain. Young age at menarche was also associated with low back pain. One theory is that increased estrogen results in increased laxity of joints and ligaments. This increased laxity then leads to LBP. Like results from other studies, association was found between the number of children and low back pain.^{12,13} A previous population-based survey found a linear association between the number of live births and chronic low back pain. Results from our study suggest that there is association between the parity and low back pain. In a population-based survey among women younger age at first pregnancy was also associated with a high prevalence of ever having low back pain.¹³ Our study shows association between younger age at first pregnancy and low back pain (Table 3). In 1995, Brynhildsen et al reported that many health professionals believe that there is an association between oral contraceptive uses with low back pain, despite the lack of scientific evidence.¹⁴ Unlike this study our study showed no association between oral contraceptive use and LBP. We found menarche at young age (<11 YEARS) to be not associated with LBP. Although LBP is positively correlated with menstruation in women and back pain is a common symptom of the premenstrual syndrome, there are only few studies describing the association between the menstrual pattern and musculoskeletal disorders.^{12,15}

CONCLUSION

Hormonal and reproductive factors like an irregular or prolonged menstrual cycle are associated with low back pain, suggesting that these factors are associated with musculoskeletal pain in general. Factors related to increased estrogen levels like past pregnancy, young maternal age at first birth, duration of oral contraceptive use may specifically increase the risk of low back pain. More research is needed to examine these associations and unravel biologic explanations.

REFERENCES

1. Waddell G. Volvo award in clinical sciences. A new clinical model for the treatment of low back pain. *Spine*. 1987;12(7):632-44.
2. Orvieto R, Achiron A, Ben-Rafael Z, et al. Low back pain of pregnancy. *Acta Obstet Gynecol Scand*. 1994;73:209-14.
3. Kristiansson P, Svardsudd K, von Schoultz B. Back pain during pregnancy: A prospective study. *Spine*. 1996;21:702-9.
4. Wang SM, Dezinno P, Maranets I, et al. Low back pain during pregnancy: Prevalence, risk factors and outcomes. *Obstet Gynecol*. 2004;104:65-70.

5. Morgen IM, Pohjanen AI. Low back pain and pelvic pain during pregnancy. *Spine* 2005;30:983-91.
6. Balague F, Troussier B, Salminen JJ. Non-specific low back pain in children and adolescents: risk factors. *Eur Spine J.* 1999;8(6):429-38.
7. Ebbehoj NE, Hansen FR, Harreby MS, Lassen CF. [Low back pain in children and adolescents. Prevalence, risk factors and prevention] *Ugeskr Laeger.* 2002;164(6):755-8.
8. Van Tulder MW, Koes B, Bombardier C. Low back pain. *Best Pract Res Clin Rheumatol.* 2002;16(5):761-75.
9. Hestback L, Leboeuf-Yds C, Manniche C. Low back pain: What is the long term course? A review of studies of general patient populations. *Eur Spine J.* 2003;12(2):149-65.
10. Wijnhoven HA, HCW ds Vet, HA Smit-Spinae. 2006;31(13):1496-502.
11. Budeiri DJ, Li Wan Po A, Dornan JC. Clinical trails of treatments of premenstrual syndrome: Entry criteria and scales for measuring treatment outcomes. *Br J Obstet Gynaecol.* 1994;101:689-95.
12. Svensson HO, Andersson GB, Hagstad A, et al. The relationship of low back pain to pregnancy and gynaecologic factors. *Spine.* 1990;15:371-5.
13. Silman AJ, Ferry S, Papageorgion AC et al. Number of children as a risk factor for low back pain in men and women. *Arthritis Rheum.* 1995;38:1232-5.
14. Brynhildsen J, Ekblad S, Hammar M. Oral contraceptives and low back pain. Attitudes among physicians, mid wives and physiotherapists. *Acta Obstet Gynecol Scand.* 1995;74:714-7.
15. Brynhildsen JO, Bjors E, Skarsgard C, et al. Is hormone replacement therapy a risk factor for low back pain among postmenopausal women? *Spine.* 1998;23:809-13.

