

Treatment of Post-traumatic Bone Defects with infection in Long Bones

Hari Om Agarwal¹, Harjit K. Singh Chawla², Ashish Garg³, Girish Sahni¹

¹Associate Professor, Deptt of Orthopaedics, Govt Medical College, Patiala.

²Assistant Professor, Deptt of Orthopaedics, Govt Medical College, Patiala

³Senior Resident, Deptt of Orthopaedics, All India Institute of Medical Sciences, Bathinda

ABSTRACT

Background: Open fractures are a challenging condition to treat because they are frequently compounded by infection and nonunion. Traditional bone defect care strategies are mostly focused on fracture union rather than infection prevention. The goal of this study is to use the Masquelet approach to examine the outcome of a post-traumatic defect with infection in long bones. This method is a two-step process. Stage I surgery includes debridement and the placement of an antibacterial spacer in the bone defect. Stage II surgery involved removing the spacer while preserving the induced membrane that had grown on the spacer's surface and filling the bone-gap with morselized iliac crest bone-graft within the membrane sleeve.

Methods: There were 22 patients in this study (18 males and 4 females), all of them had infected long bone fractures with a bone defect. The average length of the bone defect was 3.5 centimetres. The duration of follow-up varied from 6 to 15 months.

Results: After an average of 11.5 weeks following the first step of surgery, radiological union was achieved. After stage 1, no patient had any remaining infection. After radiological union, all of the patients were able to mobilise with full weight bearing and a satisfactory range of motion in the adjoining joints.

Conclusion: With favourable outcomes, this treatment can be used on infected fractures with bone loss on a regular basis. Antibiotic cement spacers, used in conjunction with complete debridement, minimise the risk of infection. The graft is revascularized through induced biomembrane. In most circumstances, union may be predicted; nonetheless, the length of time it takes to reach an agreement is a constraint. The technique is low-cost and does not necessitate any additional training or equipment. Despite the fact that it is a two-stage procedure, it does not necessitate several surgeries as in traditional approaches.

Keywords: Antibiotic cement spacer, induced biomembrane bone grafting, management of bone defect, management of compound fractures, Masquelet technique

Article History

Received: 28.11.2021

Accepted: 10.12.2021

*Corresponding Author

Dr. Girish Sahni

Associate Professor, Deptt of Orthopaedics, Govt Medical College, Patiala.

Email: dr.hariom@yahoo.com

Copyright: © the author(s). IABCR is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.




This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial

INTRODUCTION

The tibia is one of the most commonly fractured long bones.¹ Open fractures are more common in the tibia than in any other major long bone.¹ Incidences of complex open injuries of the limbs are on the rise owing to the increased number of high energy vehicular accidents in recent times, which subsequently giving rise to more cases of infected non-unions. High-energy fractures may be associated with bone loss, compartment syndrome, neurovascular injury and infection. Segmental bone defects resulting from traumatic injuries are complicated problems with significant long-term morbidity.² Historically, due to the difficulty in managing segmental long bone defects, amputation was the preferred

treatment. Various researchers over the years have used many different approaches to deal with these complex problems.³ But, it has not been possible to address all the problems mentioned above by using any single technique. In 1986, A.C. Masquelet conceived and developed an original reconstruction technique for large diaphyseal bone defects, based on the notion of the induced membrane.⁴⁻⁶ The principle of Masquelet technique involves provoking a reaction to a foreign body by placing a cement spacer in the bone defect. The membrane induced by this foreign body is in fact a biological chamber which prevents graft resorption

Access this article online	
Website: www.iabcr.org	Quick Response code
DOI: 10.21276/iabcr.2022.8.1.6	

How to cite this article: Agarwal HO, Chawla HKS, Garg A, Sahni G. Treatment of Post-traumatic Bone Defects with infection in Long Bones. Int Arch BioMed Clin Res. 2022;8(1):OT1-OT3.

Source of Support: Nil, **Conflict of Interest:** None

by providing vascularization and growth factors, as shown by various clinical, experimental and basic studies.⁴⁻¹³

The present study was performed on patients of bone defect in long bones alongwith infection treated by the Masquelet technique. The aim of the present study was to analyze the results of this study and compare them with those in the literature.

METHODS

This study includes 22 post-traumatic long bone defects with infection who were treated in Department of Orthopaedics, Govt. Medical College, Patiala between July,2016 to June,2019 by Masquelet technique. The present study included 18 men and 4 women (mean age 35.5 years). In all cases, the initial trauma was an open fracture. Tibia was involved in maximum number of cases (63.63%), followed by Femur bone in 27.27% and Humerus in 9.09% cases. The mean delay between the accident and treatment of bone defects was 3.63 months. Infection was present in 100% cases. Bone defects were 5cm or larger than 5 cm in 60% of the cases. The patients were evaluated for injury type, location, soft tissue condition, length of bone defect, antibiotic used, and duration of cementation. Moreover, the type of fixation, presence of infection, and current state of all patients was recorded.

Surgical Technique

Masquelet technique is a two-step procedure (Fig. 1): First step included excision of infected or non-viable tissue, if necessary a cover flap is created. The length, alignment, and rotation of the injured limb was obtained comparable to other normal extremity. Method of fixation depended on the fracture type and location. For open fracture, with significant defect, external fixator was used temporarily (Figure 2) and then the cement spacer was placed in the bone defect following debridement and careful stabilization of the fracture fragments. We preferred to use 4 gram Vancomycin in bone polymethylmethacrylate (PMMA) while preparing bone cement spacer (Figure 2). The second step was performed at least 4 to 6 weeks after the first step and included removing the spacer and filling the biological space which has been created with small morsels of cancellous graft, 1–2 mm³, harvested from the inner cortex of iliac crest. (Figure 3). It may be associated with bone substitutes. This step was performed after any remaining infection has healed. The fracture was approached through the previous incision only.



Figure 1: AP (a) and lateral (b) radiographs of an open fracture right distal femur Gustilo Type IIIA at admission. It was initially debrided, stabilized with an external fixator



Figure 2: AP (a) and lateral (b) radiographs showing fixation with external fixator and screws and placement of antibiotic cement spacer into the defect after the wound had been adequately debrided.



Figure 3: AP (a) and lateral (b) fluoroscopic images of another case Gustilo Type IIIA distal femur showing the fracture stabilized with distal femoral locking plate after removal of cement spacer and the defect filled with cancellous autograft harvested from iliac crest.

RESULTS

A total of 22 patients were identified within the time period. The series included 18 men and 4 women, with a mean age of 35.5 years (18-52). The bone defects were located at tibia (14 cases) with involvement of the fibula, the femur (6 cases), the humerus (2 cases). Bone defects were 5cm or larger in size in 60% of the cases. Evaluation of the size of bone defect was based on radiographic parameters: length of the defect on an AP view of the lateral & medial cortices and lateral views of anterior & posterior cortices.

Two cases were closed fracture but complicated with infection or nonunion. The other cases were open fractures with bone loss (Gustilo Classification Type II or IIIA). The length of bone defect ranged from 3.5 cm to 8 cm. Primary stabilization of the fracture was ensured by external fixation in all the cases. The antibiotics used for cement spacer was vancomycin. The mean interval between the first-stage and second-stage surgeries was 40 days (30–178). All affected limbs were fixed with screws and plates. All patients showed radiological callus formation over the defect after treatment. Mean time of appearance of callus was 11.5 weeks (6-16 weeks). Consolidation of callus occurred in duration of 12-30 weeks. Non-union of long bones united in all cases except one case where complication of breakage of distal femoral locking plate occurred after 5 months of reconstruction surgery. In this case, revision surgery was done with good surgical outcome. The delay between the initial trauma and first stage of the surgical treatment of bone defect was a

mean 3.63 months (1-6 months). All the patients had infection when treatment of bone defect began. Six skin flap covers were performed before actual bone reconstruction.

DISCUSSION

Treatment of large segmental bone defects can be challenging for orthopaedic surgeons. Masquelet et al⁷ described a procedure combining induced membranes and cancellous autografts. Bone grafting of these defects is often delayed after primary fixation to allow soft tissue healing, decrease the risk of infection, and prevent graft resorption.⁸ In the present study, we have treated 22 patients of infected nonunion of long bones with bone gap by using Masquelet technique. It has the advantage of being simple, although has to be technically performed. The two-stage procedure is an advantage in case of infection because the aim of the first step is to cure infection by using antibiotic impregnated cement spacers and restore the envelope of soft tissue. Repeated debridement may be necessary. In traumatic wounds, antibiotic impregnated cement beads or spacers are often used for local antibiotic administration to the soft tissue bed. In addition, the advantages of inserting such a spacer include maintaining a well-defined void to allow for later placement of graft, providing structural support, offloading the implant, and inducing the formation of a biomembrane. Masquelet and Begue proposed that this membrane prevents graft resorption and improves vascularity and corticalization. It has been described that, after the initial placement of the antibiotic impregnated spacer, an interval of 4 to 6 weeks is needed for development and maturation of a biologically active membrane that is suitable for grafting. The spacer also maintains the defect and inhibits fibrous ingrowth.⁶

Recent literature has shown that this biomembrane can be 0.5 to 1 mm thick⁹ and has been described as both hyper-vascular and impermeable.¹⁰ Viateau et al.¹¹ studied this technique in a sheep model and found that the membrane alone was inadequate to heal a large defect. But when autologous bone graft was placed within the membrane, all the defects went on to heal.

Aho and his colleagues¹² found that the one-month-old membrane has higher osteogenesis-improving capabilities compared to two-month-old membrane; they concluded that optimal time for performing second-stage surgery may be within a month after implantation of foreign material.¹² In our series, the mean interval between the first and second surgeries is 40 days, which is comparable to other studies. Pelissier et al.¹⁰ reported that the induced membranes secrete growth factors including vascular and osteoinductive factors and could stimulate bone regeneration.

Accadbled et al.¹⁴ reported a case using a cage and nail construct, resulting in successful eradication of methicillin-resistant staphylococcus aureus infection and reconstitution of a 17 cm diaphyseal defect in the tibia.¹⁵ Apard et al.¹⁷ reported a series of 12 patients who presented with 6 cm segmental defects in the tibia, all of whom were initially fixed with an intramedullary nail. They reported healing following the second-stage procedure in 11 of 12 patients at an average of 4 months.

The technique as described by Masquelet and Begue⁶ relied on the placement of morselized cancellous autograft harvested from the iliac crests within the biomembrane lined defect. If this amount is not sufficient, demineralized allograft is added to the autograft in a ratio that does not exceed 1:3.

In our study, we used autograft harvested mainly from iliac crest, without any allograft. Biau et al.¹³ used both iliac crest corticocancellous autograft and a medial tibial cortical strut autograft to fill their large defect. Schmidmaier et al.¹⁸ described the use of cancellous autograft from the femoral canal. They showed that levels of many growth factors (fibroblast growth factor- α , platelet derived growth factor, insulin-like growth factor 1, TGF-1, and BMP-2) in femoral cancellous bone are present in higher concentrations than they are in iliac crest and platelet preparations in our study, we used Masquelet technique to treat infected post-traumatic bone defect successfully.

CONCLUSION

In conclusion, successful reconstruction of extensive bone defects is possible with the induced membrane technique of Masquelet. The technique of delayed bone grafting after initial placement of a cement spacer provides a reasonable alternative for the challenging problem of significant bone loss with infection in long bones. The bioactivity of the membrane created by filling large bony defects with cement leads to a favourable environment for bone formation and osseous consolidation of a large void. The technique is simple and advantageous in cases of primary infection also.

REFERENCES

1. A. Paige Whittle, "Fractures of the Lower Extremity," In: S. Terry Canale and H. B. James, Eds., *Campbell's Operative Orthopaedics*, Mosby, Philadelphia, 2008, pp. 3117-3146. [Citation Time(s):2]
2. J. T. Watson, M. Anders, and B. R. Moed, "Management strategies for bone loss in tibial shaft fractures," *Clinical Orthopaedics and Related Research*, no. 315, pp. 138-152, 1995
3. H. Weinberg, V. G. Roth, G. C. Robin, and Y. Floman, "Early fibular bypass procedures (tibiofibular synostosis) for massive bone loss in war injuries," *Journal of Trauma*, vol. 19, no. 3, pp. 177-181, 1979.
4. R. Hertel, A. Gerber, U. Schlegel, J. Cordey, P. Rueggsegger, and B. A. Rahn, "Cancellous bone graft for skeletal reconstruction: muscular versus periosteal bed—preliminary report," *Injury*, vol. 25, supplement 1, pp. A59-A70, 1994.
5. A. C. Masquelet, "Muscle reconstruction in reconstructive surgery: soft tissue repair and long bone reconstruction," *Langenbeck's Archives of Surgery*, vol. 388, no. 5, pp. 344-346, 2003.
6. A. C. Masquelet and T. Begue, "The concept of induced membrane for reconstruction of long bone defects," *Orthopedic Clinics of North America*, vol. 41, no. 1, pp. 27-37, 2010.
7. A. C. Masquelet, F. Fitoussi, T. Begue, and G. P. Muller, "Reconstruction of the long bones by the induced membrane and spongy autograft," *Annales de Chirurgie Plastique et Esthetique*, vol. 45, no. 3, pp. 346-353, 2000.
8. T. A. McCall, D. S. Brokaw, B. A. Jelen et al., "Treatment of large segmental bone defects with reamer-irrigator-aspirator bone graft: technique and case series," *Orthopedic Clinics of North America*, vol. 41, no. 1, pp. 63-73, 2010
9. C. Y.-L. Woon, K.-W. Chong, and M.-K. Wong, "Induced membranes—a staged technique of bone-grafting for segmental bone loss. A report of two cases and a literature review," *The Journal of Bone and Joint Surgery. American*, vol. 92, no. 1, pp. 196-201, 2010.
10. P. Pelissier, A. C. Masquelet, R. Bareille, S. M. Pelissier, and J. Amedee, "Induced membranes secrete growth factors including vascular and osteoinductive factors and could stimulate bone regeneration," *Journal of Orthopaedic Research*, vol. 22, no. 1, pp. 73-79, 2004.
11. V. Viateau, G. Guillemin, Y. Calando et al., "Induction of a barrier membrane to facilitate reconstruction of massive segmental diaphyseal bone defects: an ovine model," *Veterinary Surgery*, vol. 35, no. 5, pp. 445-452, 2006.
12. O. M. Aho, P. Lehenkari, J. Ristiniemi, S. Lehtonen, J. Risteli, and H. V. Leskela, "The mechanism of action of induced membranes in bone repair," *The Journal of Bone and Joint Surgery. American*, vol. 95, no. 7, pp. 597-604, 2013.
13. D. J. Biau, S. Pannier, A. C. Masquelet, and C. Glorion, "Case report: reconstruction of a 16-cm diaphyseal defect after Ewing's resection in a child," *Clinical Orthopaedics and Related Research*, vol. 467, no. 2, pp. 572-577, 2009.
14. F. Accadbled, P. Mazeau, F. Chotel, J. Cottalorda, J. Sales de Gauzy, and R. Kohler, "Induced-membrane femur reconstruction after resection of bone malignancies: three cases of massive graft resorption in children," *Orthopaedics & Traumatology, Surgery & Research*, vol. 99, no. 4, pp. 479-483, 2013.
15. N. T. O'Malley and S. L. Kates, "Advances on the Masquelet technique using a cage and nail construct," *Archives of Orthopaedic and Trauma Surgery*, vol. 132, no. 2, pp. 245-248, 2012.
16. L. K. Huffman, J. G. Harris, and M. Suk, "Using the bi-masquelet technique and reamer-irrigator-aspirator for post-traumatic foot reconstruction," *Foot and Ankle International*, vol. 30, no. 9, pp. 895-899, 2009.
17. T. Apard, N. Bigorre, P. Cronier, F. Duteille, P. Bizot, and P. Massin, "Two-stage reconstruction of post-traumatic segmental tibia bone loss with nailing," *Orthopaedics and Traumatology: Surgery and Research*, vol. 96, no. 5, pp. 549-553, 2010.
18. G. Schmidmaier, S. Herrmann, J. Green et al., "Quantitative assessment of growth factors in reaming aspirate, iliac crest, and platelet preparation," *Bone*, vol. 39, no. 5, pp. 1156-1163, 2006